BHUTAN

EPIDEMIC ESTIMATES

New HIV infections			
New HIV infections (all ages)	<200	<100	<100
	[<100- <500]	[<100– <200]	[<100-<500]
New HIV infections (0-14)			
	[]	[]	[]
New HIV infections (women, 15+)	<100	<100	<100
	[<100– <100]	[<100- <100]	[<100-<100]
New HIV infections (men, 15+)	<100	<100	<100
	[<100– <200]	[<100- <200]	[<100–<200]
HIV incidence per 1000 population	0.17 [0.08–0.38]	0.13 [0.05–0.28]	0.11 [0.04–0.3]
AIDS-related deaths			
AIDS-related deaths (all ages)	<100	<100	<100
	[<100- <500]	[<100– <500]	[<100- <200]
AIDS-related deaths (0-14)			
	[]	[]	[]
AIDS-related deaths (women, 15+)	<100	<100	<100
	[<100– <100]	[<100– <100]	[<100–<100]
AIDS-related deaths (men, 15+)	<100	<100	<100
	[<100– <200]	[<100- <200]	[<100– <200]
People living with HIV			
People living with HIV (all ages)	1300	1300	1300
	[590–2700]	[670–2700]	[700–2700]
People living with HIV (0–14)			
	[]	[]	[]
People living with HIV (women, 15+)	<500	<500	<500
	[<200–670]	[<200–700]	[<500–750]
People living with HIV (men, 15+)	950	920	900
	[<500–2000]	[<500–1900]	[<500–1900]
HIV prevalence (15–49)	0.3 [0.1–0.6]	0.3 [0.1–0.6]	0.3 [0.1–0.5]

LAWS AND POLICIES

Laws criminalizing the transmission of, nondisclosure of or exposure to HIV transmission

Criminalization of sex work among consenting adults

Drug use or possession for personal use is an

Criminalization of transgender people

Criminalization of same-sex sexual acts

Laws or policies restricting the entry, stay and residence of people living with HIV

Parental consent for adolescents to access HIV testing

Spousal consent for married women to access sexual and reproductive health services

Mandatory HIV testing for marriage, work or residence permits or for certain groups

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent

VIOLENCE

Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months


EXPENDITURES


Financing sources						
Last available report:						


Yes, imprisonment (up to 14 years)


offence

EPIDEMIC TRANSITION METRICS









Change in new **HIV** infections since 2010


related deaths since 2010

Incidence: prevalence

KEY POPULATIONS

Estimated size of population	 	 •••	
HIV prevalence	 	 •••	
Know their HIV status	 	 •••	
Antiretroviral therapy coverage	 	 	
Condom use	 	 	
Coverage of HIV prevention programmes	 	 	
Avoidance of health care because of stigma and discrimination	 	 	
Expenditures (0)	 		

HIV TESTING AND TREATMENT CASCADE



All ages	47% [25->95%]	37% [20–78%] 480	% [%]
Children (0-14)	% [–%]	% [%]	% [%]
Women (15+)	73% [41– >95%]	57% [32- >95%] 220	% [%]
Men (15+)	35% [18–75%]	26% [13–57%] 230	% [–%]

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

	2010	2018
Percentage of pregnant women living with HIV	%	%
accessing antiretroviral medicines	[%]	[%]
Early infant diagnasis	%	%
Early infant diagnosis	[%]	[%]

HIV COMORBIDITIES

Estimated number of incident tuberculosis cases among people living with HIV (2017)	6 [4–9]
People living with HIV who started TB preventive therapy (2017)	
Cervical cancer screening of women living with HIV	
People coinfected with HIV and hepatitis B virus receiving combined treatment	
People coinfected with HIV and hepatitis C virus starting hepatitis C treatment	

HIV PREVENTION

Adults aged 15+ years with unsuppressed

Knowledge of HIV prevention among young people aged 15-24 years

— Women — Men

Condom use at last sex with a non-marital, non-cohabiting partner

> — Women — Men

Women aged 15-49 years who have their demand for family planning satisfied by modern methods

Men aged 15–49 years who are circumcised	Not applicable
Voluntary medical male circumcisions performed according to national standards	Not applicable
People who received PrEP at least once during the reporting period	

Harm reduction

- Use of sterile injecting equipment at last injection
- Needles and syringes distributed per person who injects
- Coverage of opioid substitution therapy
- Naloxone available (2019)
- Safe injection rooms available (2019)