PAKISTAN

EPIDEMIC ESTIMATES

New HIV infections			
New HIV infections (all ages)	14 000	18 000	22 000
	[12 000–15 000]	[17 000–20 000]	[20 000–24 000]
New HIV infections (0-14)	760	1200	1400
	[640–900]	[990–1300]	[1200–1600]
New HIV infections (women, 15+)	3800	4900	5900
	[3200–4300]	[4400–5400]	[5200–6500]
New HIV infections (men, 15+)	9400	12 000	15 000
	[8300–10 000]	[11 000–14 000]	[13 000–16 000]
HIV incidence per 1000 population	0.08 [0.07–0.09]	0.1 [0.09–0.11]	0.11 [0.1–0.12]
AIDS-related deaths			
AIDS-related deaths (all ages)	1400	4700	6400
	[870–1900]	[3700–5800]	[5200–7600]
AIDS-related deaths (0–14)	<500	660	800
	[<500– <500]	[560–760]	[690–920]
AIDS-related deaths (women, 15+)	<500	1200	1800
	[<200– <500]	[930–1500]	[1500–2200]
AIDS-related deaths (men, 15+)	700	2800	3800
	[<500–1000]	[2200–3500]	[3100–4600]
People living with HIV			
People living with HIV (all ages)	67 000	120 000	160 000
	[57 000–76 000]	[110 000–140 000]	[140 000–190 000]
People living with HIV (0-14)	1800	4000	5500
	[1500–2000]	[3400–4500]	[4700–6300]
People living with HIV (women, 15+)	19 000	37 000	48 000
	[17 000–22 000]	[32 000–42 000]	[42 000–54 000]
People living with HIV (men, 15+)	46 000	84 000	110 000
	[39 000–52 000]	[73 000–96 000]	[97 000–120 000]
HIV prevalence (15–49)	<0.1 [<0.1– <0.1]	0.1 [<0.1–0.1]	0.1 [0.1–0.2]

LAWS AND POLICIES

LAWS AND POLICIES	
Laws criminalizing the transmission of, non-disclosure of or exposure to HIV transmission	No
Criminalization of sex work among consenting adults	Any criminalization or punitive regulation of sex work
Criminalization of same-sex sexual acts	Yes, death penalty
Drug use or possession for personal use is an offence	No
Criminalization of transgender people	
Laws or policies restricting the entry, stay and residence of people living with HIV	No
Parental consent for adolescents to access HIV testing	Yes, for adolescents younger than 18 years
Spousal consent for married women to access sexual and reproductive health services	
Mandatory HIV testing for marriage, work or residence permits or for certain groups	No

STIGMA AND DISCRIMINATION

Percentage of women and men aged 15–49 years who report discriminatory attitudes towards people living with HIV

2013

49

Percentage of people living with HIV denied health services because of their HIV status in the last 12 months

Percentage of people living with HIV who reported a health-care professional told others about their HIV status without their consent

VIOLENCE


Proportion of ever-married or partnered women aged 15–49 years who experienced physical or sexual violence from a male intimate partner in the past 12 months 2013 2018


14.5


EXPENDITURES


Financing sources						
			International: PEPFAR	International: Global Fund	International: all others	
Last available report: 2013		\$3 631 968		\$5 311 <i>22</i> 5	\$313 844	\$9 993 630

EPIDEMIC TRANSITION METRICS









Change in new HIV infections = 57 since 2010

57% Change in A related dear since 2010


os. s = 369% Incidence: prevalence ratio

13%

KEY POPULATIONS

Estimated size of population					
HIV prevalence	3.8%	3.7%	21.0%	5.5%	
Know their HIV status	45.0%	44.7%	39.3%	69.1%	
Antiretroviral therapy coverage					
Condom use	35.1%	22.4%	15.3%	24.4%	
Coverage of HIV prevention programmes	0.7%	1.2%	1.6%	1.1%	
Avoidance of health care because of stigma and discrimination					
Expenditures (0)					

HIV TESTING AND TREATMENT CASCADE



All ages	14% [13–16%]	10% [9–11%] 15 800	% [%]
Children (0-14)	18% [15–21%]	11% [10–13%] 610	% [%]
Women (15+)	8% [7–9%]	7% [6–8%] 3300	% [%]
Men (15+)	16% [15–19%]	11% [10–12%] 11 900	% [%]

ELIMINATION OF MOTHER-TO-CHILD TRANSMISSION

	2010	2018
Percentage of pregnant women living with HIV	1%	10%
accessing antiretroviral medicines	[1–2%]	[8–12%]
Early infant diagnosis	1.3%	1.9%
Larry Illiant diagnosis	[1.0-1.6%]	[1.6-2.2%]

HIV COMORBIDITIES

Estimated number of incident tuberculosis cases among people living with HIV (2017)	7200 [3600 –12 000]
People living with HIV who started TB preventive therapy (2017)	
Cervical cancer screening of women living with HIV	
People coinfected with HIV and hepatitis B virus receiving combined treatment	
People coinfected with HIV and hepatitis C virus starting hepatitis C treatment	

HIV PREVENTION

Adults aged 15+ years with unsuppressed viral load

Knowledge of HIV prevention among young people aged 15–24 years (2017)

Condom use at last say with a non-marital	
— Men	5.9%
— Women	1.8%

Condom use at last sex with a non-marital, non-cohabiting partner

• .	
— Women	
— Men	

Women aged 15–49 years who have their demand for family planning satisfied by modern methods

Men aged 15–49 years who are circumcised	applicable	
Voluntary medical male circumcisions performed according to national standards	Not applicable	
People who received PrEP at least once during the reporting period		

Harm reduction

 Use of sterile injecting equipment at last injection (2016) 	72.5%
 Needles and syringes distributed per person who injects (2018) 	46
 Coverage of opioid substitution therapy 	
— Naloxone available (2019)	No

- Safe injection rooms available (2019)