Pakistan

Population 2018		212 million
		Rate
Estimates of TB burden ^o , 2018	Number (thousands)	(per 100 000 population)
Total TB incidence	562 (399-754)	265 (188–355)
HIV-positive TB incidence	3.8 (2.5–5.4)	1.8 (1.2–2.5)
MDR/RR-TB incidence ⁰⁰	28 (18–40)	13 (8.4–19)
HIV-negative TB mortality	43 (35–52)	20 (16–25)
HIV-positive TB mortality	1.3 (0.83–1.8)	0.6 (0.39–0.86)

Estimated proportion of TB cases with MDR/RR-TB, 2018	
New cases	4.2% (3.2–5.3)
Previously treated cases	16% (15–17)

TB case notifications, 2018	
Total new and relapse	360 472
- % tested with rapid diagnostics at time of diagnosis	22%
- % with known HIV status	20%
- % pulmonary	80%
- % bacteriologically confirmed ⁰⁰⁰	48%
- % children aged 0-14 years	13%
- % women	42%
- % men	45%
Total cases notified	369 548
Universal health coverage and social protection	
TB treatment coverage (notified/estimated incidence), 2018	64% (48–90)
TB treatment coverage (notified/estimated incidence), 2018	64% (48–9

I B patients facing catastrophic total costs		
TB case fatality ratio (estimated mortality/estimated incidence), 2018	8%	(5–11)
TD// IIV / sere in new and relates TD patients 2010	Numero	(0/)
TB/HIV care in new and relapse TB patients, 2018	Number	(%)
Patients with known HIV status who are HIV-positive	636	<1%
- on antiretroviral therapy	417	66%
Drug-resistant TB care, 2018		
% of bacteriologically confirmed TB cases tested for rifampicin resistance $^{\circ\circ\circ}$		
- New cases		45%

	1070
- Previously treated cases	79%
Laboratory-confirmed cases*	MDR/RR-TB: 3 824, XDR-TB: 95
Patients started on treatment* **	MDR/RR-TB: 3 106, XDR-TB: 71
MDR/RR-TB cases tested for resistance to second-line drugs	2 893

93%	358 730
	220/20
79%	9673
64%	2 804
35%	77
	64% 35%

TB preventive treatment, 2018

% of HIV-positive people (newly enrolled in care) on preventive treatment	
% of children (aged < 5) household contacts of bacteriologically-confirmed	5.7% (5.2–6.3)
TB cases on preventive treatment	

```
TB financing, 2019
```

National TB budget (US\$ millions)	
Funding source: 3% domestic, 31% international, 66% unfunded	

° Ranges represent uncertainty intervals

^{oo} MDR is TB resistant to rifampicin and isoniazid; RR is TB resistant to rifampicin


ooo Calculated for pulmonary cases only

* Includes cases with unknown previous TB treatment history

** Includes patients diagnosed before 2018 and patients who were not laboratory-confirmed


Tuberculosis profile


(Rate per 100 000 population per year)




HIV-positive TB incidence

(Rate per 100 000 population per year)






Notified cases by age group and sex, 2018



Treatment success rate (%)





New and relapse

135

Retreatment, excluding relapse

- HIV-positive - MDR/RR-TB - XDR-TB





Funded domestically